

Class 09: Computing with Symbols and Numbers

Held: Tuesday, 5 February 2008

Summary: We explore some of Scheme's standard numeric functions.

Related Pages:

- EBoard.
- Lab: Numeric Computation.
- Reading: Numeric Values & Symbolic Values.

Notes:

- Are there questions on Assignment 3?
- Advance notice: Extra credit for attending Thursday's convo.
- There is only one readings for tomorrow, Representing Images as Lists of Spots
- The Mac version of DrFu is now available, but experimental. Send comments!

Overview:

- Types.
- Kinds of Numbers.
- Key Numeric Operations.

Types

- As you may have noted in your first experiments with Scheme, Scheme assigns *types* to variables.
- For example, a value might be a number, or a string, or an image identifier, or a procedure, or a drawing, or
- Computer scientists often think of types in two different ways:
 - *Data-driven*: A type is a set of values.
 - *Purpose-driven*: A type provides information on the valid operations that may be applied to a piece of data.
- We will alternate between the two definitions.
- Many languages (particularly the ones you've reported being familiar with) require you to assign a type to a variable when you declare that variable.
- Scheme does not require you to assign types of variables; it checks the type of each operand when it executes a procedure.
 - Scheme also provides procedures that let you determine the type of a value.
- As the semester progresses, you will learn new types.

Scheme's Numeric Types

- Instead of a general “numbers” type, Scheme provides a variety of kinds of numbers.
 - Integers are numbers without a fractional component.
 - Rational numbers can be expressed as the ratio of two integers.
 - Real numbers appear somewhere on the number line.
 - In mathematics, real numbers can be rational or irrational.
 - In Scheme, real numbers are all rational.
 - Complex numbers may include an *imaginary* component.
 - Our version of Scheme does not support complex numbers, but we mention them anyway.
 - You can (almost) always use an integer when a real is expected, but you cannot always use
 - Scheme also represents some numbers exactly and some numbers inexactly. (That is, it approximates some numbers.)
 - It certainly has to approximate irrational numbers.
 - But it also approximates many other numbers.
 - It may surprise you to see which numbers are represented inexactly. (We'll return to this issue later.)
 - Some important numeric predicates (procedures that return true or false): `number?`, `real?`, `integer?`, `exact?`, and `inexact?`.
-

Copyright © 2007-8 Janet Davis, Matthew Kluber, and Samuel A. Rebelsky. (Selected materials copyright by John David Stone and Henry Walker and used by permission.) This material is based upon work partially supported by the National Science Foundation under Grant No. CCLI-0633090. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation. This work is licensed under a Creative Commons Attribution-NonCommercial 2.5 License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc/2.5/> or send a letter to Creative Commons, 543 Howard Street, 5th Floor, San Francisco, California, 94105, USA.